

QUADRATIC FORMULA PHYSICS

THE QUADRATIC FORMULA IN PHYSICS: SOLVING MOTION AND MORE

QUADRATIC FORMULA PHYSICS IS A POWERFUL MATHEMATICAL TOOL THAT UNLOCKS SOLUTIONS TO MANY PROBLEMS IN MECHANICS, ELECTROMAGNETISM, AND BEYOND. WHEN YOU ENCOUNTER SITUATIONS INVOLVING PARABOLIC TRAJECTORIES, PROJECTILE MOTION, OR OSCILLATING SYSTEMS, THE QUADRATIC FORMULA OFTEN BECOMES YOUR BEST FRIEND. THIS ARTICLE WILL DEMYSTIFY THE QUADRATIC FORMULA'S APPLICATION IN PHYSICS, BREAKING DOWN ITS DERIVATION, EXPLAINING ITS CORE CONCEPTS, AND SHOWCASING ITS PRACTICAL USES. WE'LL EXPLORE HOW IT HELPS DETERMINE TIME, POSITION, AND OTHER CRUCIAL VARIABLES IN DYNAMIC PHYSICAL SCENARIOS, MAKING COMPLEX CALCULATIONS MANAGEABLE AND PROVIDING A DEEPER UNDERSTANDING OF THE UNDERLYING PHYSICAL PRINCIPLES.

TABLE OF CONTENTS

UNDERSTANDING THE QUADRATIC EQUATION

DERIVING THE QUADRATIC FORMULA

THE QUADRATIC FORMULA: A DEEPER DIVE

APPLICATIONS OF THE QUADRATIC FORMULA IN PHYSICS

PROJECTILE MOTION AND THE QUADRATIC FORMULA

KINEMATIC EQUATIONS AND QUADRATIC SOLUTIONS

OSCILLATORY MOTION AND QUADRATIC SOLUTIONS

OTHER PHYSICS SCENARIOS REQUIRING THE QUADRATIC FORMULA

COMMON PITFALLS AND BEST PRACTICES

CONCLUSION

UNDERSTANDING THE QUADRATIC EQUATION

AT ITS HEART, THE QUADRATIC FORMULA IS DESIGNED TO SOLVE A SPECIFIC TYPE OF EQUATION: THE QUADRATIC EQUATION. YOU'LL RECOGNIZE A QUADRATIC EQUATION BY ITS STANDARD FORM: $ax^2 + bx + c = 0$. HERE, ' a ', ' b ', AND ' c ' ARE COEFFICIENTS, AND ' x ' IS THE VARIABLE WE WANT TO FIND. THE KEY CHARACTERISTIC IS THAT THE HIGHEST POWER OF THE VARIABLE IS 2, HENCE THE "QUADRATIC" PART. THESE EQUATIONS ARE FUNDAMENTAL BECAUSE THEY DESCRIBE MANY NATURAL PHENOMENA, FROM THE PATH OF A THROWN BALL TO THE ELECTRICAL RESPONSE OF A CIRCUIT.

WHY IS SOLVING QUADRATIC EQUATIONS SO IMPORTANT IN PHYSICS? BECAUSE MANY PHYSICAL LAWS AND RELATIONSHIPS ARE INHERENTLY QUADRATIC. FOR INSTANCE, THE DISTANCE AN OBJECT TRAVELS UNDER CONSTANT ACCELERATION IS PROPORTIONAL TO THE SQUARE OF TIME ($d = vt + \frac{1}{2}at^2$), WHICH IS A QUADRATIC RELATIONSHIP. OR CONSIDER THE ENERGY OF A SPRING, WHICH IS PROPORTIONAL TO THE SQUARE OF ITS DISPLACEMENT ($E = \frac{1}{2}kx^2$). WITHOUT A RELIABLE WAY TO SOLVE THESE EQUATIONS, MANY CORE PHYSICS PROBLEMS WOULD REMAIN INTRACTABLE.

DERIVING THE QUADRATIC FORMULA

THE DERIVATION OF THE QUADRATIC FORMULA IS A TESTAMENT TO ALGEBRAIC INGENUITY. WE START WITH THE GENERAL QUADRATIC EQUATION, $ax^2 + bx + c = 0$, AND AIM TO ISOLATE ' x '. THE PROCESS INVOLVES COMPLETING THE SQUARE, A TECHNIQUE THAT TRANSFORMS ONE SIDE OF THE EQUATION INTO A PERFECT SQUARE TRINOMIAL. FIRST, WE DIVIDE THE ENTIRE EQUATION BY ' a ' TO GET $x^2 + (b/a)x + (c/a) = 0$. THIS STEP IS CRUCIAL TO ENSURE THE COEFFICIENT OF x^2 IS 1, WHICH IS NECESSARY FOR COMPLETING THE SQUARE.

NEXT, WE MOVE THE CONSTANT TERM TO THE RIGHT SIDE: $x^2 + (b/a)x = -c/a$. NOW COMES THE MAGIC OF COMPLETING THE SQUARE. WE TAKE HALF OF THE COEFFICIENT OF ' x ' (WHICH IS b/a), SQUARE IT, AND ADD IT TO BOTH SIDES OF THE EQUATION. HALF OF b/a IS $b/(2a)$, AND SQUARING IT GIVES US $b^2/(4a^2)$. SO, WE ADD $b^2/(4a^2)$ TO BOTH SIDES: $x^2 + (b/a)x + b^2/(4a^2) = -c/a + b^2/(4a^2)$.

THE LEFT SIDE IS NOW A PERFECT SQUARE: $(x + b/(2a))^2$. ON THE RIGHT SIDE, WE FIND A COMMON DENOMINATOR TO COMBINE THE TERMS: $(x + b/(2a))^2 = (b^2 - 4ac) / (4a^2)$. TAKING THE SQUARE ROOT OF BOTH SIDES, WE GET $x + b/(2a) = \pm \sqrt{(b^2 - 4ac) / (4a^2)}$. FINALLY, ISOLATING 'x' BY SUBTRACTING $b/(2a)$ FROM BOTH SIDES YIELDS THE FAMOUS QUADRATIC FORMULA: $x = [-b \pm \sqrt{(b^2 - 4ac)}] / (2a)$.

THE QUADRATIC FORMULA: A DEEPER DIVE

THE QUADRATIC FORMULA, $x = [-b \pm \sqrt{(b^2 - 4ac)}] / (2a)$, IS MORE THAN JUST A SET OF SYMBOLS; IT'S A UNIVERSAL KEY TO FINDING THE ROOTS OF ANY QUADRATIC EQUATION. THE BEAUTY OF THIS FORMULA LIES IN ITS ABILITY TO PROVIDE UP TO TWO DISTINCT SOLUTIONS FOR 'x', REPRESENTED BY THE ' $\pm$ ' SIGN. THESE TWO SOLUTIONS OFTEN CORRESPOND TO DIFFERENT PHYSICAL SCENARIOS OR TIME POINTS IN A PROBLEM.

LET'S BREAK DOWN THE COMPONENTS. THE TERM UNDER THE SQUARE ROOT, $b^2 - 4ac$, IS KNOWN AS THE DISCRIMINANT. THE DISCRIMINANT IS INCREDIBLY INSIGHTFUL BECAUSE ITS VALUE TELLS US ABOUT THE NATURE OF THE ROOTS WITHOUT ACTUALLY CALCULATING THEM. IF THE DISCRIMINANT IS POSITIVE, THERE ARE TWO DISTINCT REAL ROOTS, INDICATING TWO POSSIBLE VALID SOLUTIONS IN MANY PHYSICS PROBLEMS. IF THE DISCRIMINANT IS ZERO, THERE IS EXACTLY ONE REAL ROOT (A REPEATED ROOT), MEANING THERE'S A UNIQUE SOLUTION. IF THE DISCRIMINANT IS NEGATIVE, THERE ARE NO REAL ROOTS, WHICH IMPLIES THAT THE PHYSICAL SITUATION DESCRIBED BY THE QUADRATIC EQUATION IS NOT POSSIBLE UNDER THE GIVEN CONDITIONS.

APPLICATIONS OF THE QUADRATIC FORMULA IN PHYSICS

THE QUADRATIC FORMULA IS NOT JUST AN ABSTRACT MATHEMATICAL CONCEPT; IT'S A WORKHORSE IN PRACTICAL PHYSICS. WHENEVER YOU ENCOUNTER A PROBLEM WHERE A PHYSICAL QUANTITY DEPENDS ON THE SQUARE OF ANOTHER, OR WHEN YOU'RE TRYING TO FIND THE TIME IT TAKES FOR SOMETHING TO HAPPEN, THE QUADRATIC FORMULA IS LIKELY TO BE INVOLVED. ITS ABILITY TO HANDLE SECOND-ORDER RELATIONSHIPS MAKES IT INDISPENSABLE FOR ANALYZING MOTION, ENERGY, AND WAVE PHENOMENA.

ONE OF THE MOST COMMON AREAS WHERE THE QUADRATIC FORMULA SHINES IS IN KINEMATICS, THE STUDY OF MOTION. MANY KINEMATIC EQUATIONS, WHICH DESCRIBE HOW OBJECTS MOVE, ARE QUADRATIC IN NATURE. BY PLUGGING THE KNOWN PHYSICAL PARAMETERS INTO THESE EQUATIONS AND REARRANGING THEM INTO THE STANDARD QUADRATIC FORM, WE CAN USE THE FORMULA TO SOLVE FOR UNKNOWN LIKE TIME OR DISPLACEMENT. THIS ALLOWS PHYSICISTS AND ENGINEERS TO PREDICT THE BEHAVIOR OF OBJECTS WITH REMARKABLE ACCURACY.

PROJECTILE MOTION AND THE QUADRATIC FORMULA

PROJECTILE MOTION IS A CLASSIC EXAMPLE WHERE THE QUADRATIC FORMULA IS INDISPENSABLE. IMAGINE LAUNCHING A BALL INTO THE AIR. ITS TRAJECTORY FOLLOWS A PARABOLIC PATH, A SHAPE DIRECTLY LINKED TO QUADRATIC EQUATIONS. THE VERTICAL COMPONENT OF MOTION IS GOVERNED BY GRAVITY, WHICH CAUSES A CONSTANT DOWNWARD ACCELERATION. THE EQUATION DESCRIBING THE VERTICAL POSITION (y) OF THE PROJECTILE AT ANY TIME (t) TYPICALLY LOOKS LIKE $y = y_0 + v_{0y}t + \frac{1}{2}at^2$, WHERE y_0 IS THE INITIAL HEIGHT, v_{0y} IS THE INITIAL VERTICAL VELOCITY, AND 'a' IS THE ACCELERATION DUE TO GRAVITY (USUALLY -9.8 m/s^2).

IF WE WANT TO FIND OUT WHEN THE PROJECTILE REACHES A CERTAIN HEIGHT, SAY $y = 0$ (WHEN IT HITS THE GROUND), WE SET THE EQUATION TO 0: $0 = y_0 + v_{0y}t + \frac{1}{2}at^2$. REARRANGING THIS INTO THE STANDARD $ax^2 + bx + c = 0$ FORM, WE GET $\frac{1}{2}at^2 + v_{0y}t + y_0 = 0$. HERE, THE VARIABLE 't' IS WHAT WE ARE SOLVING FOR. APPLYING THE QUADRATIC FORMULA WITH $a = \frac{1}{2}a$, $b = v_{0y}$, AND $c = y_0$ ALLOWS US TO CALCULATE THE SPECIFIC TIME(S) AT WHICH THE PROJECTILE IS AT THAT HEIGHT. OFTEN, ONE SOLUTION REPRESENTS THE TIME GOING UP, AND THE OTHER THE TIME COMING DOWN, OR ONE MIGHT BE A PHYSICALLY IRRELEVANT NEGATIVE TIME.

KINEMATIC EQUATIONS AND QUADRATIC SOLUTIONS

BEYOND JUST PROJECTILE MOTION, OTHER KINEMATIC EQUATIONS FREQUENTLY LEAD TO QUADRATIC SOLUTIONS. FOR INSTANCE, CONSIDER THE EQUATION: $\Delta x = v_0 t + \frac{1}{2} a t^2$. THIS EQUATION RELATES DISPLACEMENT (Δx) TO INITIAL VELOCITY (v_0), TIME (t), AND ACCELERATION (a). IF WE KNOW THE DISPLACEMENT AND INITIAL VELOCITY AND WANT TO FIND THE TIME IT TOOK TO ACHIEVE THAT DISPLACEMENT UNDER CONSTANT ACCELERATION, WE REARRANGE THIS EQUATION TO FIT THE QUADRATIC FORM.

LET'S SAY WE'RE INTERESTED IN FINDING THE TIME ' t ' WHEN AN OBJECT TRAVELING WITH AN INITIAL VELOCITY v_0 AND CONSTANT ACCELERATION ' a ' REACHES A DISPLACEMENT Δx . WE WOULD REWRITE THE EQUATION AS: $\frac{1}{2} a t^2 + v_0 t - \Delta x = 0$. IN THIS SCENARIO, OUR ' a ' IN THE QUADRATIC FORMULA IS $\frac{1}{2} a$, OUR ' b ' IS v_0 , AND OUR ' c ' IS $-\Delta x$. SOLVING THIS USING THE QUADRATIC FORMULA GIVES US THE SPECIFIC TIME(S) AT WHICH THAT DISPLACEMENT OCCURS. THIS IS INCREDIBLY USEFUL FOR ANALYZING CAR ACCELERATION, THE MOTION OF FALLING OBJECTS, AND MANY OTHER EVERYDAY PHYSICAL SCENARIOS.

OSCILLATORY MOTION AND QUADRATIC SOLUTIONS

OSCILLATORY MOTION, LIKE THAT OF A PENDULUM OR A MASS ON A SPRING, OFTEN INVOLVES SECOND-ORDER DIFFERENTIAL EQUATIONS, WHICH, WHEN SOLVED OR ANALYZED IN CERTAIN CONTEXTS, CAN LEAD BACK TO QUADRATIC EQUATIONS. FOR EXAMPLE, IN ANALYZING THE ENERGY STORED IN A SPRING, THE POTENTIAL ENERGY (PE) IS GIVEN BY $PE = \frac{1}{2} k x^2$, WHERE ' k ' IS THE SPRING CONSTANT AND ' x ' IS THE DISPLACEMENT FROM EQUILIBRIUM. IF YOU'RE ASKED TO FIND THE DISPLACEMENT ' x ' WHEN THE SPRING HAS A SPECIFIC AMOUNT OF POTENTIAL ENERGY, YOU WOULD SET UP THE EQUATION:

$$\frac{1}{2} k x^2 = PE$$

REARRANGING THIS TO THE STANDARD FORM, WE GET:

$$\frac{1}{2} k x^2 - PE = 0$$

HERE, $a = \frac{1}{2} k$, $b = 0$, AND $c = -PE$. APPLYING THE QUADRATIC FORMULA, $x = [0 \pm \sqrt{0^2 - 4 (\frac{1}{2} k) (-PE)}] / (2 \cdot \frac{1}{2} k) = \pm \sqrt{2k PE} / k$. THIS SHOWS THAT THE DISPLACEMENT IS DIRECTLY RELATED TO THE STORED ENERGY, AND THE QUADRATIC FORMULA PROVIDES A CLEAR PATHWAY TO CALCULATE IT.

OTHER PHYSICS SCENARIOS REQUIRING THE QUADRATIC FORMULA

THE REACH OF THE QUADRATIC FORMULA EXTENDS BEYOND BASIC MECHANICS. IN ELECTRICAL CIRCUITS, FOR INSTANCE, YOU MIGHT ENCOUNTER EQUATIONS DESCRIBING THE CHARGE OR CURRENT OVER TIME THAT ARE QUADRATIC. FOR EXAMPLE, THE DISCHARGE OF A CAPACITOR THROUGH A RESISTOR IN AN RC CIRCUIT CAN INVOLVE SECOND-ORDER DIFFERENTIAL EQUATIONS, AND SOLVING FOR TIME CONSTANTS OR SPECIFIC CHARGE VALUES CAN LEAD TO QUADRATIC SOLUTIONS. SIMILARLY, IN OPTICS, PROBLEMS INVOLVING FOCAL LENGTHS AND IMAGE FORMATION USING LENS EQUATIONS SOMETIMES REQUIRE SOLVING QUADRATIC EQUATIONS TO DETERMINE THE POSITION OF AN OBJECT OR IMAGE.

EVEN IN THERMODYNAMICS, CERTAIN RELATIONSHIPS MIGHT EXHIBIT QUADRATIC BEHAVIOR. THE KINETIC THEORY OF GASES, FOR EXAMPLE, DEALS WITH THE DISTRIBUTION OF MOLECULAR SPEEDS, AND WHILE THE MAXWELL-BOLTZMANN DISTRIBUTION ITSELF ISN'T A SIMPLE QUADRATIC EQUATION, INTERMEDIATE CALCULATIONS OR SPECIFIC PROBLEM FORMULATIONS WITHIN THIS DOMAIN MIGHT NECESSITATE ITS USE. ESSENTIALLY, ANY TIME A PHYSICAL LAW OR A DERIVED EQUATION RESULTS IN A TERM SQUARED, BE PREPARED TO POTENTIALLY CALL UPON THE QUADRATIC FORMULA.

COMMON PITFALLS AND BEST PRACTICES

WHILE THE QUADRATIC FORMULA IS A POWERFUL TOOL, IT'S IMPORTANT TO USE IT CORRECTLY TO AVOID ERRORS. ONE COMMON PITFALL IS MISIDENTIFYING THE COEFFICIENTS ' a ', ' b ', AND ' c ' WHEN REARRANGING AN EQUATION INTO THE STANDARD FORM. ALWAYS ENSURE THAT THE EQUATION IS SET EQUAL TO ZERO AND THAT THE TERMS ARE IN THE CORRECT ORDER. PAY

CLOSE ATTENTION TO THE SIGNS OF THE COEFFICIENTS; A MISPLACED NEGATIVE SIGN CAN DRAMATICALLY ALTER THE RESULT.

ANOTHER FREQUENT MISTAKE INVOLVES THE DISCRIMINANT, $b^2 - 4ac$. IF THE DISCRIMINANT IS NEGATIVE, IT MEANS THERE ARE NO REAL SOLUTIONS. IN A PHYSICS CONTEXT, THIS USUALLY SIGNIFIES THAT THE SCENARIO DESCRIBED BY THE EQUATION IS PHYSICALLY IMPOSSIBLE UNDER THE GIVEN CONDITIONS. DON'T TRY TO FORCE A REAL NUMBER SOLUTION WHERE NONE EXISTS. FURTHERMORE, WHEN INTERPRETING THE TWO SOLUTIONS PROVIDED BY THE ' $\pm$ ' SIGN, ALWAYS CONSIDER THE PHYSICAL CONTEXT. ONE SOLUTION MIGHT BE POSITIVE AND PHYSICALLY MEANINGFUL, WHILE THE OTHER MIGHT BE NEGATIVE AND IRRELEVANT, OR VICE VERSA, DEPENDING ON THE PROBLEM.

HERE ARE SOME BEST PRACTICES TO KEEP IN MIND:

- ALWAYS REWRITE YOUR PHYSICS EQUATION IN THE STANDARD QUADRATIC FORM: $ax^2 + bx + c = 0$.
- CAREFULLY IDENTIFY YOUR ' a ', ' b ', AND ' c ' COEFFICIENTS, INCLUDING THEIR SIGNS.
- WHEN CALCULATING THE DISCRIMINANT ($b^2 - 4ac$), BE METICULOUS WITH YOUR ARITHMETIC.
- UNDERSTAND WHAT THE DISCRIMINANT TELLS YOU ABOUT THE NATURE OF THE ROOTS.
- ALWAYS CHECK THE PHYSICAL REASONABLENESS OF YOUR SOLUTIONS. DOES A NEGATIVE TIME MAKE SENSE? DOES A DISTANCE GREATER THAN THE EARTH'S CIRCUMFERENCE MAKE SENSE FOR A THROWN BALL?
- USE A CALCULATOR WITH A DEDICATED QUADRATIC SOLVER FUNCTION IF AVAILABLE, BUT UNDERSTAND THE UNDERLYING FORMULA.
- DOUBLE-CHECK YOUR SUBSTITUTION INTO THE FORMULA ITSELF.

BY ADHERING TO THESE PRACTICES, YOU CAN SIGNIFICANTLY IMPROVE THE ACCURACY AND EFFICIENCY OF YOUR PROBLEM-SOLVING IN PHYSICS. THE QUADRATIC FORMULA IS A FRIEND, BUT LIKE ANY TOOL, IT REQUIRES CAREFUL HANDLING TO YIELD THE BEST RESULTS.

THE JOURNEY THROUGH UNDERSTANDING AND APPLYING THE QUADRATIC FORMULA IN PHYSICS REVEALS ITS PROFOUND SIGNIFICANCE. FROM THE SIMPLE ARC OF A PROJECTILE TO THE COMPLEXITIES OF OSCILLATORY SYSTEMS, THIS MATHEMATICAL BEDROCK PROVIDES ESSENTIAL SOLUTIONS. ITS ABILITY TO NAVIGATE SECOND-ORDER RELATIONSHIPS ALLOWS US TO QUANTIFY AND PREDICT THE BEHAVIOR OF THE PHYSICAL WORLD WITH IMPRESSIVE PRECISION. MASTERING ITS APPLICATION OPENS DOORS TO A DEEPER COMPREHENSION OF FUNDAMENTAL PHYSICS PRINCIPLES AND EMPOWERS YOU TO TACKLE A WIDER ARRAY OF CHALLENGING PROBLEMS WITH CONFIDENCE AND CLARITY.

FAQ

Q: WHEN DO I TYPICALLY NEED TO USE THE QUADRATIC FORMULA IN PHYSICS?

A: YOU MOST COMMONLY ENCOUNTER THE NEED FOR THE QUADRATIC FORMULA IN PHYSICS WHEN DEALING WITH MOTION UNDER CONSTANT ACCELERATION (KINEMATICS), PROJECTILE MOTION, ENERGY CALCULATIONS INVOLVING SQUARED QUANTITIES (LIKE SPRING POTENTIAL ENERGY OR KINETIC ENERGY), AND CERTAIN ASPECTS OF OSCILLATORY OR WAVE MOTION WHERE SECOND-ORDER RELATIONSHIPS ARISE.

Q: WHAT DOES THE DISCRIMINANT ($b^2 - 4ac$) TELL ME IN A PHYSICS PROBLEM?

A: THE DISCRIMINANT IS A CRUCIAL INDICATOR OF THE PHYSICAL REALITY OF YOUR SOLUTION. A POSITIVE DISCRIMINANT MEANS THERE ARE TWO DISTINCT REAL SOLUTIONS, OFTEN REPRESENTING TWO DIFFERENT TIMES OR POSITIONS WHERE A CONDITION IS

MET. A ZERO DISCRIMINANT MEANS THERE'S A SINGLE, UNIQUE REAL SOLUTION. A NEGATIVE DISCRIMINANT INDICATES THAT THE PHYSICAL SITUATION DESCRIBED BY THE EQUATION IS IMPOSSIBLE UNDER THE GIVEN PARAMETERS, MEANING THERE ARE NO REAL SOLUTIONS.

Q: CAN THE QUADRATIC FORMULA GIVE ME NEGATIVE ANSWERS IN PHYSICS PROBLEMS, AND ARE THEY VALID?

A: YES, THE QUADRATIC FORMULA CAN YIELD NEGATIVE ANSWERS. HOWEVER, WHETHER A NEGATIVE ANSWER IS VALID DEPENDS ENTIRELY ON THE PHYSICAL CONTEXT. FOR INSTANCE, A NEGATIVE TIME IS USUALLY UNPHYSICAL IF YOU'RE MEASURING TIME FROM A STARTING POINT IN THE PAST, BUT IT MIGHT BE MATHEMATICALLY VALID IF YOUR REFERENCE FRAME IS SET DIFFERENTLY. SIMILARLY, NEGATIVE DISPLACEMENT OR VELOCITY MIGHT BE PERFECTLY VALID DEPENDING ON THE DIRECTION DEFINED AS POSITIVE.

Q: HOW DO I ENSURE I'VE CORRECTLY IDENTIFIED 'A', 'B', AND 'C' FROM A PHYSICS EQUATION?

A: THE BEST WAY IS TO REARRANGE YOUR PHYSICS EQUATION INTO THE STANDARD QUADRATIC FORM, $ax^2 + bx + c = 0$, WHERE 'X' IS THE VARIABLE YOU'RE SOLVING FOR (OFTEN TIME 'T', POSITION 'X', ETC.). ENSURE THE EQUATION IS SET EQUAL TO ZERO. THEN, 'A' IS THE COEFFICIENT OF THE SQUARED TERM, 'B' IS THE COEFFICIENT OF THE LINEAR TERM, AND 'C' IS THE CONSTANT TERM. PAY VERY CLOSE ATTENTION TO THE SIGNS OF THESE COEFFICIENTS.

Q: WHAT HAPPENS IF I GET TWO POSITIVE SOLUTIONS FROM THE QUADRATIC FORMULA IN A PHYSICS PROBLEM?

A: IF YOU GET TWO POSITIVE SOLUTIONS, IT TYPICALLY MEANS THAT THE PHYSICAL CONDITION YOU'RE INVESTIGATING IS MET AT TWO DISTINCT, POSITIVE POINTS IN TIME OR AT TWO DISTINCT POSITIONS. FOR EXAMPLE, IN PROJECTILE MOTION, AN OBJECT MIGHT REACH A CERTAIN HEIGHT TWICE: ONCE ON ITS WAY UP AND ONCE ON ITS WAY DOWN. YOU'LL NEED TO ANALYZE THE CONTEXT TO DETERMINE WHICH SOLUTION, OR IF BOTH, ARE RELEVANT TO YOUR SPECIFIC QUESTION.

[Quadratic Formula Physics](#)

Quadratic Formula Physics

Related Articles

- [quantum physics weird facts](#)
- [principal axis in physics](#)
- [physics tutor nyc](#)

[Back to Home](#)